

POST SURGERY CARE INSTRUCTIONS: Upper Lip Lift

*These instructions apply to most individuals. We may modify our recommendations to you, depending on your specific needs. Please follow these instructions UNLESS we notify you in writing to do something different.

WHAT TO EXPECT AFTER SURGERY

SWELLING:

- Swelling is a normal part of the healing process and typically peaks 72 hours post-op.
- A cold compress or ice may be used to ease discomfort and swelling, if desired, around the surgical area.
- DO NOT place cold compresses or ice on the incisions. DO NOT place ice directly on the skin- place a towel or cloth between skin and ice.
- You will notice a gradual reduction in swelling after 2 weeks.
- By week 4, expect a noticeable improvement in appearance of the surgical area. Residual swelling typically resolves in 6-8 weeks.

BRUISING:

- You may experience some bruising a few days after surgery- this is normal.
- A cold compress or ice may be used to ease discomfort and bruising, if desired, around the surgical area.
- DO NOT place cold compresses or ice on the incisions.
- You will notice a gradual reduction in bruising after 2 weeks.
- By week 3-4, expect a noticeable improvement in appearance of the surgical area as your body heals.

SAFETY / CAREGIVER

After surgery, you will be weak and drowsy. You will need a responsible ADULT caregiver to stay with you 24 hours after surgery to monitor your health and support you.

MEDICATION SAFETY

Dr. Camp uses a multimodal pain medication regimen to help keep pain tolerable and our patients comfortable. Stay ahead of the pain for the first 24 hours by following the schedule of medications outlined on your medication chart. After 48 hours, Dr Camp recommends avoiding the use of narcotics like hydrocodone (norco) and muscle relaxers such as valium. Continue taking Celecoxib (Celebrex) and Gabapentin every 12hours to maintain comfort. **Constant, severe, worsening pain should be reported to our office immediately.**

It is important to take the medications as prescribed. DO NOT drive while taking narcotics, muscle relaxers or any medication that causes drowsiness. Additionally, DO NOT drink alcohol while still taking any of the prescription medications.

POSITION AND ACTIVITY

HEAD POSITION: Keep your head elevated for 3 weeks to reduce swelling. When resting or sleeping, lie on your back with your head elevated 30-45 degrees. We recommend using a wedge pillow, an adjustable bed, or a recliner to maintain this position. You may lay flat after 3 weeks. DO NOT lay on your side or stomach for 6 weeks.

FACIAL MOVEMENT: Please minimize facial movements involving the surgical area as much as possible for the first few weeks.

ACTIVITY:

WEEKS 1 - 3

- For the first 3 weeks post-op, walk as much as you would like indoors.

- Avoid raising your heart rate over 100 or increasing your blood pressure.
- No sweating

WEEKS 3 – 6

- You may ease back into light daily activities and hygiene as you normally would. Listen to your body and avoid stretching or motions that cause too much pain. Your normal range of motion will return as you heal.
- You may increase activity with faster walking.
- No "exercise" or high impact activity (biking, elliptical, weights or resistance training)
- Avoid vigorous household activities like vacuuming, house cleaning, etc.
- Do not lift, push, or pull anything greater than 10lbs.

After 6 WEEKS

- Expect your energy to return to normal 6-8 weeks after surgery.
- Ease back into physical activity, starting light and increasing frequency and intensity gradually over 3-6 months, depending on your health, physical stamina and pre-operative level of fitness.
- There is no weight limit, per se, but listen to your body. Discomfort is your body telling you to back off.

SUN EXPOSURE: Avoid sun exposure for 6 weeks. Treated areas are more likely to sunburn for up to 1 year. Wear sunscreen daily once your sutures are removed.

DRIVING: Avoid driving until your vision is clear and unobstructed and no longer taking narcotic pain medication.

DEEP BREATHING

Practicing deep breathing after surgery will help strengthen and clear your lungs, lowering the risk of postoperative pneumonia. Inhale through your nose and slowly exhale through your mouth.

NUTRITION

Adequate fluid and nutritional intake are CRITICAL for proper healing. You may not feel hungry after surgery. It is essential that you spread your required fluid and nutritional intake evenly throughout the day. Protein intake is EXTREMELY important for wound healing. Start with a liquid diet and then progress to a soft foods diet.

Eat soft foods for the first week after surgery, such as soup, pasta, yogurt, protein shakes, smoothies, and Jello. Avoid foods that are difficult to chew that will disrupt the incision- including hard breads, meats, burgers, pizza, tacos, etc. After 1 week, you may eat a regular diet. Limit carbonated beverages and spicy foods that can cause gas and bloating. Avoid salty foods which will increase swelling.

SHOWERING AND HYGIENE

- For patients that have a LIP LIFT WITHOUT ADDITIONAL SURGICAL PROCEDURES, you may shower **24 hours after surgery**. If you have a LIP LIFT WITH ADDITIONAL SURGICAL PROCEDURES, please follow the showering and hygiene instructions of the other procedures.
- Wash your face during the first shower using the facial cleanser provided.
- Be careful pulling clothing over your head.
- Resume skin care regimen after the sutures are removed.
- DO NOT submerge the surgical area in water for at least 6 weeks, until the incisions are healed.
- DO NOT wear lip products (lipstick, gloss, Chapstick, balm, etc.) for 6 weeks. Aquaphor is OK to keep your lips hydrated.

SUTURES AND INCISION CARE

- You may have small amounts of drainage around the incision sites the first few days after surgery.
- Apply the prescribed Erythromycin ointment to incision sites twice daily, using a clean Q-tip.
- Sutures are removed in the office 7-10 days post-op.
- DO NOT apply bandages or other materials to the surgical site unless otherwise instructed.

BLOOD CLOT PREVENTION

It is essential that during the day you get up EVERY HOUR and walk around indoors for 10 minutes while you recover to reduce the risk of developing blood clots after surgery.

CONSTIPATION PROTOCOL

You may not have a bowel movement for several days after surgery, this is normal. Constipation can be prevented by staying hydrated, moving, and taking MiraLAX 4x/day in a liquid of your choice. MiraLAX does not cause cramping and is very safe/effective. Diarrhea can be caused by antibiotics taken postoperatively- please notify us if you experience this.

URINATION

We want you to urinate within 4 hours of surgery. Ideally, you should urinate at least twice before you go to bed the evening of surgery. If you are not able to urinate, please notify us ASAP.

NURSE PHONECALL

Your post-op nurse will call you the day after your surgery. Please answer so we can make sure you are healing normally and answer any questions you may have that are not covered in the post-op instructions. We want to speak to you. We are also happy to speak with your caregiver, but we want to hear your voice and have you answer our post-op questions. Please write down all your questions and save your questions for this call. Our RNs have scheduled time to check on our immediate post-op patients.

QUESTIONS AFTER SURGERY

Steven Camp MD Plastic Surgery

Phone: (817) 228-4315

Text: (833) 784-3316

Office Hours: Monday – Friday, 9 am – 5 pm

For post-op questions, please call or text the office and leave a message for our post-op nurse. Please note there may be delays in getting back to you with non-urgent questions as our RNs are in clinic treating patients during the week. Social media should NOT be used to contact us regarding your care. ***Prior to calling, please review your post-op instructions to ensure your question is not already answered.***

EMERGENCIES & AFTER-HOURS

NONURGENT: For nonurgent and routine post-op questions after business hours, please TEXT OR CALL THE OFFICE AND LEAVE A VOICEMAIL. We will contact you within 24 hours regarding non-urgent matters.

NONURGENT issues include:

- Medication refill requests
- Routine post-op questions
- Appointment scheduling

URGENT: For urgent, non-life-threatening medical issues or concerns, please CALL OUR OFFICE IMMEDIATELY to be connected to our after-hours emergency line. The after-hours emergency line rings directly to Dr. Steven Camp and Sara Camp AG-ACNP, APRN, MSN. If they do not answer, please leave a detailed voice message. They are most likely on another call and will contact you immediately.

URGENT issues include:

- Surgical drain issues
- Fever of 101 or greater
- Allergic reaction to medications
- Increasing swelling or redness around the incision
- Drainage from incision that is yellow or green
- Drainage from incision that has a foul odor
- Severe pain that is not relieved by medications
- Severe nausea or vomiting that is not relieved by medications

EMERGENCY: If you are experiencing a life-threatening emergency, CALL 911 OR GO TO THE NEAREST EMERGENCY ROOM IMMEDIATELY. While we cannot provide immediate assistance, we ask that you update us once you are safe.

EMERGENT issues include:

- Shortness of breath or difficulty breathing
- Chest pain
- Loss of consciousness
- Calf pain or tenderness, lower leg/calf swelling, one leg more swollen than the other • Uncontrolled bleeding

COMMON QUESTIONS

Jewelry/Piercings: If you have piercings, you can have non-metal spacers placed before you arrive at the surgery center, as long as the piercing area is a NON-SURGICAL area. Otherwise, not even a spacer is allowed. Please don't call us and ask about this as there are no exceptions. It's for your protection and keeps the tools used in surgery from burning your skin!

Menstrual Cycle: What if I'm on my period the day of surgery? This will NOT hinder you from having surgery. Please wear a pad (not a tampon) to surgery. You will be provided with pads and disposable underwear at the surgery center.

Exercise: See 'Activity Section' for details. Doing so earlier can mess up your results and prolong healing.

SEX: Refrain from sex for at least 4-6 weeks after surgery. Resume with caution and avoid pressure or rubbing on surgical sites.

Swimming/Baths: AVOID for at least 6 weeks after surgery. All your incisions must be COMPLETELY healed and closed before you do this. Otherwise, you could get a bad infection! This includes lakes, oceans, hot tub, pool, and bath water.

Scar Treatments: A few weeks after your procedure and when incisions completely closed, our nurse will direct you to start using Silicone Scar Strips or silicone scar gel- apply with clean hands. Scar Away is a good over the counter brand. You can get these products at CVS, Walgreens, Wal-Mart, or on Amazon. You may start applying scar strips for 12 hrs. on, then 12 hrs. off every day. Scar gel may be applied every 12 hours. Continue using for several months after surgery.

Post-Operative Travel: Typically, we recommend not traveling for at least 6 weeks for vacations! Necessary travel before this time should be done cautiously, taking care to adhere to lifting and activity restrictions to avoid complications. This is the timeframe in which you have the highest risk of getting an infection or an incision opening up.

Alcohol: Avoid for 2 weeks before surgery and at least 2 weeks after surgery. You must be off all of your medications for at least ONE full week before you drink alcohol. Alcohol is NOT good for healing!

Smoking: No smoking or second-hand exposure to smoke for AT LEAST 6 weeks after surgery.

Tanning: Avoid excessive sun exposure 10 days prior to surgery. If you have a sunburn, surgery may have to be delayed. Avoid direct sun exposure of surgical site skin for at least 6 weeks after surgery. Sunblock with UVA/ UVB protection SPF 35 or greater is recommended to protect skin with any sun exposure. Note: bathing suits and thin clothing offer minimal sun protection. You should always apply sunblock under such garments.

Spray Tan: Avoid for 2 weeks prior to surgery and for 6 weeks after surgery.

Tattoos: Avoid for 6 weeks before surgery and 3 months after surgery to decrease the risk of infection.

Pets: No pets sleeping in your bed for 6 weeks. Do not walk large dogs with a leash for 6 weeks after surgery.

Nails: Do not replace dip, gel, or acrylic nails until cleared by your nurse.

Should I search online or ask my friends if I have any questions about my procedure or recovery? You know the answer to this one! Google "experts," reading social media "expert groups" or asking your friend that had plastic surgery causes a lot of problems and unnecessary anxiety. Please call us if you have any questions or concerns. Dr. Camp is a board-certified plastic surgeon with nearly 10 years in private practice. It is dangerous to attempt to diagnose yourself or depend on the advice of bloggers, social media sites or friends and family. We are here to help, ask us anything!